



DEPARTMENT OF
PLANT AND SOIL SCIENCES

2020 Oklahoma Replicated Agronomic Cotton Evaluation (RACE) Trial Report



cotton.okstate.edu

ntokcotton.org

Contributors

Seth Byrd – Cotton Extension Specialist

Bradley Wilson – Graduate Research Assistant

Cayden Catlin – Graduate Research Assistant

Andrea Althoff – Graduate Research Assistant

Jerry Goodson – Extension Assistant, IPM

Cameron Murley – Superintendent, OPREC

Aaron Henson – Tillman Co. Extension Educator

Josh Bushong – Area Agronomist

Brian Pugh – Area Agronomist and District Program Leader

Greg Hartman – Washita and Beckham counties Extension Educator

Danny Cook – Roger Mills Co. Extension Educator

Cody Linker – Lincoln Co. Extension Educator

Gary Strickland – Greer and Jackson Co. Extension Educator and Dryland Cropping Systems Specialist

Acknowledgements

First and foremost, we would like to thank the Oklahoma cotton producers who serve as cooperators and allow us to use their land, equipment, and time to conduct these trials. These trials wouldn't be possible without their cooperation and support. Financial support for these trials was provided by the Cotton Inc. Oklahoma State Support Committee and the participating seed companies. We would like to thank Khawar Arain, Brendan Kelly, and the staff of the Texas Tech University Fiber and Biopolymer Research Institute for providing fiber classing services for these trials. And finally, we would like to thank the Oklahoma Cotton Council for their support of both the cotton agronomy program at Oklahoma State University and the entire Oklahoma cotton industry.

Participating Seed Companies

Americot/NexGen

Deltapine

Dyna-Gro Seed

FiberMax/Stoneville

PhytoGen Cottonseed

2020 Season Overview

Planting conditions were generally favorable for producers in 2020, although rain events in May delayed planting on a large proportion of acreage to late May and early June. In fact, the southwest and west central locations of our 2020 RACE trials were all planted between May 29th and June 3rd. Moisture was rapidly diminishing by the end of the first week of June and hot, dry conditions dominated the primary cotton regions of the state through the remainder of the month and persisted into early July. Various significant rain events were received from mid-July into early August, although there were variations on how much and when rain fell across certain areas. For example, while the Oklahoma Mesonet site located in Elk City recorded 3.4" of rain during July, stations in Tipton and Altus each recorded 1.4". However, this was reversed in August, with only 0.85" of rain received at Elk City (over half occurring over the last 3 days of the month), compared to 2.1" at both Altus and Tipton. This July through August rain was beneficial for both dryland and irrigated cotton. It coincided with the squaring through early/peak bloom growth stages for the majority of cotton in Oklahoma which helped to reduce water stress during these critical growth stages and optimize yield.

There was an overall favorable outlook as the crop entered September, although the dry August had certainly lowered the optimism once surrounding the dryland crop. Then a cold snap struck western Oklahoma on September 9 – 10th, with daytime highs struggling to reach 60 at best, and nighttime lows flirting with freezing in many areas. Although this 2-day cold snap was followed by a return to warmer temperatures, it is likely partially to blame for the struggles many had with defoliation and boll opening later in the year. It is more likely that this short-term cold snap was an extreme sample of a larger issue, a cooler than average September with total heat unit accumulation falling 100 - 150 below average. It is likely that the slower leaf drop and boll opening, as well as the variability in micronaire or low micronaire that was common to many areas in 2020 is to blame on the weather conditions the crop experienced throughout September. There was one final hurdle to cross as an ice storm struck the area in late October, coating open bolls in ice and resulting in lint stringing out of the bolls. Luckily, there was little wind with this system or in subsequent weather events, and the actual loss of lint due to fallout was surprisingly low given the severity of the storm and the number of open bolls present on the plants. The overall feeling from 2020 was that while both yields and quality were good, maybe even great in some cases, they generally fell short of expectations that had built by the favorable conditions throughout the first three months of the season.

A total of 25 commercial varieties and one experimental variety from five seed companies were evaluated across five locations in the 2020 RACE Trials (Table 1). There were a multitude of insect and herbicide traits represented within these varieties, including both two and three gene Bt and herbicide traits (Table 2). Seasonal temperature patterns from the southwest and panhandle regions are illustrated by the monthly heat unit accumulation for 2020 compared to the 12-year average for Altus (Fig. 1) and Goodwell (Fig. 2). Dryland performance pooled across our southwest region dryland trials is presented in Table 3, while variety performance compared to the trial average of entries that were included in at least four locations is presented in Table 4. Since there are variations in variety entries across the various regions of our state, these tables provide a hint of variety performance stability for areas un-represented by our current RACE trial locations.

Table 1. Seed company participants and variety abbreviations for entries in 2020 Oklahoma RACE Trials.

Seed Company	Variety Entries	Abbreviation
Deltapine®	1646 B2XF	DP 1646 B2XF
	1820 B3XF	DP 1820 B3XF
	1822 XF	DP 1822 XF
	1948 B3XF	DP 1948 B3XF
	2012 B3XF	DP 2012 B3XF
	2020 B3XF	DP 2020 B3XF
	2038 B3XF	DP 2038 B3XF
	2044 B3XF	DP 2044 B3XF
Dyna-Gro®	3317 B3XF	DG 3317 B3XF
	3385 B2XF	DG 3385 B2XF
FiberMax®	1621 GL	FM 1621 GL
	2398 GLTP	FM 2398 GLTP
NexGen®	2982 B3XF	NG 2982 B3XF
	3729 B2XF	NG 3729 B2XF
	3930 B3XF	NG 3930 B3XF
	4098 B3XF	NG 4098 B3XF
	4936 B3XF	NG 4936 B3XF
	5711 B3XF	NG 5711 B3XF
PhytoGen®	350 W3FE	PHY 350 W3FE
	400 W3FE	PHY 400 W3FE
	480 W3FE	PHY 480 W3FE
Stoneville®	4990 B3XF	ST 4990 B3XF
	5600 B2XF	ST 5600 B2XF
	5610 B3XF	ST 5610 B3XF
	5707 B2XF	ST 5707 B2XF

Table 2. Insect and herbicide trait glossary for 2020 Oklahoma RACE Trial entries.

Insect Trait	Abbreviation	Bt Proteins
Bollgard II®	B2	Cry1Ac + Cry2Ab
Bollgard 3®	B3	Cry1Ac + Cry2Ab + Vip3A
TwinLink®	T	Cry1Ab + Cry2Ae
TwinLink Plus®	TP	Cry1Ab + Cry2Ae + Vip3Aa19
Widestrike 3®	W3	Cry1Ac + Cry1F + Vip3A
Herbicide Trait	Abbreviation	Herbicide Tolerances
FlexEnlist®	FE	Glyphosate + Glufosinate + 2,4-D
GlyTol LibertyLink®	GL	Glyphosate + Glufosinate
XtendFlex®	XF	Glyphosate + Glufosinate + Dicamba

Monthly Heat Unit Accumulation, Altus, OK




Figure 1. Monthly heat unit (DD 60) accumulation from Altus, OK Mesonet station.

Monthly Heat Unit Accumulation Goodwell, OK




Figure 2. Monthly heat unit (DD 60) accumulation from Goodwell, OK Mesonet station.



DEPARTMENT OF
PLANT AND SOIL SCIENCES

Table 3. 2020 Southwest Dryland Pooled RACE Trial Results.

Variety	Average Lint Yield ¹ lbs./acre	Yield Deviation ²
NG 4098 B3XF	877 a	+9.5%
ST 5600 B2XF	837 a	+3.0%
NG 3930 B3XF	835 a	+3.8%
ST 5707 B2XF	826 a	+1.8%
DP 1646 B2XF	823 a	+1.0%
DP 1948 B3XF	802 ab	-1.1%
NG 5711 B3XF	738 b	-10.1%
NG 4936 B3XF	732 b	-10.9%

¹Varieties included across Greer, Jackson, and Tillman trial locations.

²Percent difference between yield of variety and trial yield average across all three locations

Table 4. 2020 Variety Yield Deviation¹ (minimum 4 locations).

Variety	Greer Dry	Jackson Dry	Jackson Irr.	Lincoln Dry	Texas Dry	Texas Irr.	Tillman Dry	Tillman Irr.	Washita Dry	Average
NG 4098 B3XF	+7.46%	+2.37%	-2.87%	+8.55%			+13.88%	-1.58%	+19.63%	+6.78%
NG 3930 B3XF	+2.67%	-1.10%		+3.46%	+4.60%	+2.56%	+6.58%	+2.30%	+2.34%	+2.93%
DP 1948 B3XF	-1.15%	-1.52%	+9.86%				-0.87%	+5.07%		+2.28%
ST 5707 B2XF	+0.30%	+1.54%	-4.71%	-3.61%	-1.04%		+3.38%	+4.34%	+2.50%	+0.34%
ST 5600 B2XF	+3.88%	+3.95%	+4.73%		-25.55%		+1.86%	-5.72%	+8.32%	-1.22%
NG 5711 B3XF	-4.0%	-3.87%	+1.52%				-18.25%	+0.27%	-1.42%	-4.29%
DP 2020 B3XF			-9.38%	-1.86%		-10.52%	+7.29%	-1.32%	-16.29%	-5.34%
NG 4936 B3XF	-9.51%	-4.90%	-8.17%	-12.27%			-14.21%	-3.37%	-12.56%	-9.28%
DP 2012 B3XF				-0.52%	-14.10%	-12.86%			-11.71%	-9.80%

¹Percent increase or decrease in yield compared to the trial average



2020 Greer Co. Dryland RACE Trial

Planted: May 29th

Seeding Rate: 25,000 40" row spacing

Harvested: November 7th, stripper harvested

Table 5. Greer County dryland RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return per Acre ² (\$/acre)
NG 4098 B3XF	851 a	34.40 a	4.19 bc	1.15 b	32.9 b	82.1 b-d	54.10	413
ST 5600 B2XF	823 ab	33.79 ab	4.23 b	1.10 de	32.4 b	81.8 cd	53.63	399
NG 3930 B3XF	813 ab	34.54 a	4.89 a	1.09 e	27.3 d	82.0 cd	51.30	378
ST 5610 B3XF	809 ab	34.77 a	4.08 bc	1.11 d	30.5 c	81.9 cd	53.98	394
DP 1646 B2XF	798 ab	33.74 a-c	3.83 bc	1.16 b	29.9 c	81.4 d	52.25	371
ST 5707 B2XF	794 ab	32.30 bc	4.72 a	1.13 cd	34.5 a	82.7 a-c	53.25	390
DP 1948 B3XF	783 a-c	31.98 b-d	3.79 c	1.20 a	34.2 a	83.8 a	54.32	383
ST 4990 B3XF	771 bc	31.79 cd	4.02 bc	1.11 de	30.0 c	81.6 cd	54.03	374
NG 5711 B3XF	760 bc	32.10 b-d	3.85 bc	1.12 cd	30.9 c	82.0 cd	53.52	364
NG 4936 B3XF	717 c	30.23 d	4.00 bc	1.14 bc	29.9 c	83.3 ab	54.13	345
Average	792	32.96	4.16	1.13	31.3	82.3	53.45	381
p-value	0.0451	0.0018	0.0003	<0.0001	<0.0001	0.0158	0.4399	0.2758
<i>pLSD</i>	69	1.98	0.43	0.03	1.2	1.3	NS ³	NS
CV	7.82	5.32	10.19	3.22	1.32	7.57	3.08	9.51

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³NS; no significant statistical difference between varieties at p-value ≤ 0.1 .



2020 Jackson Co. Dryland Strip Trial

Planted: June 3rd

Seeding Rate: 20,000, 40" row spacing

Harvested: November 19th, stripper harvested

Table 6. Jackson County dryland strip trial results.

Variety	Lint Yield (lbs./acre)	Turnout %	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return per Acre ² (\$/acre)
ST 5600 B2XF	802	35.87	4.72	1.13	32.0	82.5	53.85	398
DP 1646 B2XF	799	38.48	4.73	1.16	28.6	82.3	54.55	400
NG 4098 B3XF	790	34.89	4.03	1.15	31.8	81.2	54.10	389
ST 5707 B2XF	784	33.26	4.75	1.16	34.0	83.4	55.15	405
NG 3930 B3XF	763	36.83	4.89	1.08	26.8	83.9	53.35	375
DP 1948 B3XF	760	33.10	4.01	1.16	32.3	82.4	54.10	377
NG 5711 B3XF	742	33.41	3.96	1.12	29.9	81.1	56.00	381
NG 4936 B3XF	734	32.82	4.79	1.15	30.0	84.0	56.45	380
Average	772	34.83	4.49	1.14	30.7	82.6	54.69	388

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.



DEPARTMENT OF
PLANT AND SOIL SCIENCES

2020 Jackson Co. Irrigated RACE Trial

Planted: May 29th

Seeding Rate: 42,000 38" row spacing

Irrigation: sub-surface drip located on 72" spacing in furrow

Harvested: November 12th, picker harvested

Table 7. Jackson County irrigated RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return ² (\$/acre)
DP 1948 B3XF	1,882 a	40.75 a-c	4.09	1.29 a	31.6 bc	84.1 a	54.35	951 a
PHY 400 W3FE	1,814 ab	41.34 ab	4.28	1.20 cd	31.9 b	83.4 a-d	54.55	915 ab
DP 2038 B3XF*	1,796 a-c	42.50 a	4.32	1.14 e	30.3 cd	82.3 d	54.65	912 ab
ST 5600 B2XF	1,794 a-c	41.35 ab	4.85	1.22 b-d	31.0 b-d	83.5 a-d	54.65	908 ab
PHY 480 W3FE	1,782 a-c	38.45 b-d	3.97	1.19 d	30.2 cd	83.7 a-c	53.32	877 a-c
NG 5711 B3XF	1,739 a-d	40.35 a-c	3.99	1.23 bc	30.9 b-d	82.6 b-d	53.07	851 a-c
PHY 350 W3FE	1,714 a-e	38.89 b-d	4.26	1.20 cd	31.1 b-d	82.9 a-d	54.75	866 a-c
NG 4098 B3XF	1,663 b-e	36.45 d	3.81	1.29 a	35.2 a	82.5 cd	52.10	786 c
ST 5707 B2XF	1,632 c-e	36.40 d	4.20	1.22 b-d	34.5 a	84.0 a	54.93	840 bc
FM 2398 GLTP	1,612 de	41.14 ab	4.67	1.22 b-d	31.3 b-d	83.8 ab	53.65	795 c
NG 4936 B3XF	1,573 de	37.77 cd	4.18	1.24 b	29.9 d	83.7 a-c	54.77	789 c
DP 2020 B3XF	1,552 e	35.96 cd	3.90	1.23 bc	30.3 cd	83.1 a-d	54.85	776 c
Average	1,713	39.28	4.21	1.22	31.5	83.3	54.14	856
p-value	0.007	0.001	0.1013	<0.0001	<0.0001	0.0912	0.6302	0.0261
pLSD	168	3.06	NS ³	0.03	1.4	1.3	NS	107
CV	7.65	6.76	11.38	3.48	5.62	1.03	3.09	9.11

*Grower-selected entry.

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) – seed cost per acre.

³NS; no significant statistical difference between varieties at p-value ≤0.1.



DEPARTMENT OF
PLANT AND SOIL SCIENCES

2020 Lincoln Co. Dryland RACE Trial

Planted: May 5th

Seeding Rate: 34,000 38" row spacing

Harvested: November 8th, picker harvested

Table 8. Lincoln County dryland RACE trial results.

Variety	Lint Yield ¹ (lbs./acre)	Turnout (%)	Micronaire	Length (Inches)	Strength (g/tex)	Uniformity (%)	Loan Value ² (cents/lb.)	Return per Acre ³ (\$/acre)
NG 4098 B3XF	1,223	40.92	3.83	1.20	35.7	82.5	54.28	599
PHY 350 W3FE	1,197	39.83	3.27	1.15	33.0	82.0	48.58	523
PHY 480 W3FE	1,189	40.64	3.86	1.12	31.6	82.9	54.05	584
PHY 400 W3FE	1,184	42.22	3.63	1.12	31.4	81.4	53.85	578
NG 3930 B3XF	1,166	40.35	3.48	1.18	31.8	83.3	51.92	552
DP 2012 B3XF	1,121	41.26	3.71	1.17	33.2	82.6	54.20	547
DP 2020 B3XF	1,106	40.71	3.64	1.17	31.2	82.3	54.48	541
ST 5707 B2XF	1,086	39.87	4.04	1.18	35.8	82.6	54.23	543
FM 1621 GL	1,007	44.06	4.72	1.13	34.5	82.1	54.00	505
NG 4936 B3XF	988	40.29	3.82	1.13	30.5	82.2	53.95	475
Average	1,127	41.02	3.80	1.16	32.9	82.4	53.35	545

¹Four replications planted of each variety, approximately two plots were combined into a single round module producing two modules from each variety.

²Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

³Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.



2020 Texas Co. Dryland RACE Trial

Planted: May 14th

Seeding Rate: 35,000 30" row spacing

Harvested: November 5th, stripper harvested without bur extractor

Table 9. Texas County dryland RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return ² (\$/acre)
DG 3317 B3XF	420 a	28.29 ab	2.95 a	1.04 c	26.7 bc	79.7	40.75 a	110 a
DP 1822 XF	413 a	28.92 a	3.01 a	1.12 a	28.3 ab	81.4	46.43 a	145 a
DG 3385 B2XF	406 ab	28.17 ab	3.16 a	1.05 bc	26.3 bc	80.8	43.97 a	118 a
NG 3930 B3XF	387 ab	27.03 a-c	2.93 ab	1.08 a-c	27.1 bc	81.3	44.00 a	114 a
NG 3729 B2XF	379 ab	25.78 a-c	2.94 a	1.09 ab	27.6 ab	80.1	44.45 a	114 a
ST 5707 B2XF	366 ab	25.43 a-c	2.81 ab	1.09 a-c	29.9 a	80.1	41.97 a	106 a
NG 2982 B3XF	365 a-c	24.71 bc	2.25 c	1.05 bc	27.7 ab	80.1	32.38 b	49 b
DP 2012 B3XF	318 bc	23.27 c	2.48 bc	1.06 bc	25.1 c	79.5	32.60 b	42 b
ST 5600 B2XF	275 c	24.01 c	2.95 a	1.06 bc	28.3 ab	80.2	40.37 a	53 b
Average	370	26.18	2.83	1.07	27.5	80.4	40.77	95
p-value	0.0605	0.0508	0.0154	0.0755	0.0227	0.3516	0.0016	0.0018
<i>pLSD</i>	90	3.78	0.46	0.05	2.3	NS ³	6.37	47
CV	16.92	10.21	12.54	3.41	6.29	1.30	14.27	44.26

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³NS; no significant statistical difference between varieties at p-value ≤0.1.



2020 Texas Co. Irrigated RACE Trial

Planted: May 12th

Seeding Rate: 50,000 30" row spacing

Irrigation: lateral move overhead system

Harvested: November 9th, stripper harvested

Table 10. Texas County irrigated RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb)	Return ² (\$/acre)
NG 2982 B3XF	1,447 a	29.67 a-d	2.52 cd	1.13 cd	29.27 ab	82.0	43.58 a-d	453 a
DG 3317 B3XF	1,305 ab	34.04 a	2.89 a-c	1.08 f	28.00 bc	81.4	45.58 a	481 a
DG 3385 B2XF	1,301 ab	31.58 ab	3.12 a	1.09 ef	27.43 cd	80.9	44.87 ab	507 a
NG 3729 B2XF	1,267 ab	30.06 a-c	2.94 ab	1.16 ab	27.77 b-d	81.1	37.08 de	480 a
DP 1820 B3XF	1,156 bc	30.75 a-c	2.91 a-c	1.19 a	30.63 a	80.7	35.70 e	429 a
NG 3930 B3XF	1,139 bc	28.78 b-d	2.74 a-d	1.15 bc	27.70 b-d	81.9	38.15 b-e	410 a
DP 2020 B3XF	994 c	26.14 cd	2.57 b-d	1.12 cd	25.13 e	79.6	44.45 a-c	262 b
DP 2012 B3XF	968 cd	24.82 d	2.54 b-d	1.11 de	26.00 de	80.2	42.90 a-d	275 b
ST 4990 B3XF	790 de	25.82 cd	2.40 d	1.16 b	27.93 bc	80.2	36.82 de	206 b
ST 5610 B3XF	742 e	25.88 cd	2.37 d	1.12 d	27.73 b-d	80.1	37.60 c-e	203 b
Average	1,111	28.75	2.70	1.13	27.8	80.8	40.67	371
p-value	<0.0001	0.0167	0.0120	<0.0001	0.0005	0.1094	0.0396	<0.0001
pLSD	204	4.99	0.41	0.03	1.8	NS ³	7.19	126
CV	22.74	13.57	11.77	3.15	6.15	1.39	12.79	35.79

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³NS; no significant statistical difference between varieties at p-value \leq 0.1.



2020 Tillman Co. Dryland RACE Trial

Planted: June 2nd

Seeding Rate: 25,000 40" row spacing

Harvested: November 19th, stripper harvested

Table 11. Tillman County dryland RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return ² (\$/acre)
NG 4098 B3XF	990 a	36.23 ab	4.13 a-c	1.21 a	32.7 a	81.8	54.23 a	489 a
DP 2020 B3XF	932 ab	34.86 b-d	3.95 bc	1.15 bc	28.1 d	82.3	53.65 b	456 ab
NG 3930 B3XF	926 ab	33.39 de	4.06 bc	1.15 bc	29.9 b-d	82.6	53.80 b	458 ab
ST 5707 B2XF	898 ab	34.17 c-e	4.33 ab	1.11 d	31.9 ab	82.5	53.78 b	449 ab
ST 5600 B2XF	885 b	37.24 a	4.50 a	1.13 cd	29.8 b-d	82.3	53.63 b	432 ab
DP 1646 B2XF	872 b	36.08 ab	3.82 c	1.21 a	29.5 cd	82.0	53.80 b	424 b
DP 1948 B3XF	862 b	35.08 bc	3.82 c	1.21 a	31.2 a-c	83.0	54.27 a	425 b
NG 4936 B3XF	746 c	32.82 e	3.99 bc	1.17 b	29.3 cd	83.3	53.95 ab	359 c
NG 5711 B3XF	711 c	34.16 c-e	3.83 c	1.15 bc	28.7 d	82.0	53.73 b	339 c
Average	869	34.89	4.05	1.17	30.1	82.4	53.87	426
p-value	0.0007	0.0002	0.0358	<0.0001	0.0057	0.6515	0.0048	0.0009
<i>pLSD</i>	104	1.48	0.43	0.03	2.16	<i>NS</i> ³	0.33	57
CV	11.46	4.58	7.63	3.20	5.94	1.17	0.51	12.69

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³*NS*; no significant statistical difference between varieties at p-value ≤ 0.1 .



DEPARTMENT OF
PLANT AND SOIL SCIENCES

2020 Tillman Co. Irrigated RACE Trial

Planted: June 1st

Seeding Rate: 40,000 40" row spacing

Irrigation: center pivot

Harvested: November 20th, picker harvested

Table 12. Tillman County irrigated RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return ² (\$/acre)
DP 1948 B3XF	1021	39.68 a	4.27 bc	1.21	33.2 b	82.9	54.85	492
ST 5707 B2XF	1014	36.80 b	4.36 b	1.19	35.7 a	85.0	54.40	498
NG 3930 B3XF	994	40.28 a	4.37 b	1.15	29.0 c	82.7	54.23	475
NG 5711 B3XF	975	39.44 a	4.18 bc	1.19	30.8 c	82.8	54.38	461
DP 2020 B3XF	959	36.73 b	3.81 c	1.21	29.6 c	83.4	54.28	449
NG 4098 B3XF	957	37.17 b	3.94 bc	1.24	34.3 ab	83.2	54.32	443
NG 4936 B3XF	939	36.99 b	4.18 bc	1.21	29.4 c	84.1	54.43	443
ST 5600 B2XF	917	39.00 a	4.94 a	1.18	33.1 b	84.2	53.47	422
Average	972	38.26	4.25	1.20	31.9	83.5	54.30	460
p-value	0.596	0.002	0.0095	0.1038	<0.0001	0.2728	0.2454	0.3427
<i>pLSD</i>	<i>NS</i> ³	1.18	0.49	<i>NS</i>	2.0	<i>NS</i>	<i>NS</i>	<i>NS</i>
CV	7.01	4.33	9.42	2.87	8.15	1.50	1.14	9.15

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³*NS*; no significant statistical difference between varieties at p-value ≤ 0.1 .



2020 Washita Co. Dryland RACE Trial

Planted: June 2nd

Seeding Rate: 30,000 40" row spacing

Harvested: November 11th, stripper harvested

Table 13. Washita County dryland RACE trial results.

Variety	Lint Yield (lbs./acre)	Turnout (%)	Micronaire	Length (inches)	Strength (g/tex)	Uniformity (%)	Loan Value ¹ (cents/lb.)	Return ² (\$/acre)
NG 4098 B3XF	789 a	36.65 a	4.92 cd	1.15 a	32.9 a	81.4	53.23 a	363 a
DP 2044 B3XF	746 ab	36.49 a	4.76 de	1.15 a	31.6 a	81.5	53.08 a	358 a
ST 5600 B2XF	715 a-c	36.87 a	5.36 a	1.10 bc	29.8 b	82.6	48.95 cd	299 b
ST 5707 B2XF	676 bc	34.11 bc	5.41 a	1.12 abc	32.9 a	82.7	50.37 b-d	300 b
NG 3930 B3XF	675 bc	36.54 a	5.03 bc	1.09 bc	28.9 bc	82.7	51.17 a-c	298 b
NG 5711 B3XF	651 cd	36.75 a	4.89 c-e	1.08 c	29.5 bc	81.0	52.57 ab	290 bc
NG 3729 B2XF	634 c-e	36.79 a	5.25 ab	1.08 c	27.2 d	81.4	48.33 d	258 b-d
DP 2012 B3XF	583 de	35.33 ab	4.70 de	1.09 bc	28.1 cd	81.9	51.93 ab	248 cd
NG 4936 B3XF	577 de	32.47 c	4.72 de	1.13 ab	29.7 b	83.6	52.95 a	255 b-d
DP 2020 B3XF	552 e	34.73 ab	4.65 e	1.09 bc	28.1 cd	81.9	52.92 a	239 d
Average	660	35.67	4.97	1.11	28.9	82.1	51.55	291
p-value	0.0004	0.0066	<0.0001	0.0143	<0.0001	0.1606	0.0013	0.0003
<i>pLSD</i>	90	2.23	0.26	0.05	1.5	NS ³	2.32	49
CV	13.42	5.04	6.85	3.24	6.91	1.42	4.25	17.53

¹Color and leaf grades set to base levels (41 and 4, respectively) due to lack of proper lint cleaner on gin.

²Return per acre calculated as (lint yield per acre X loan value per pound) - seed cost per acre.

³NS; no significant statistical difference between varieties at p-value ≤ 0.1 .